

Marquette University
e-Publications@Marquette

Mathematics, Statistics and Computer Science
Faculty Research and Publications

Mathematics, Statistics and Computer Science,
Department of

6-1-1991

Corrigendum to "Taxonomies of Model-theoretically Defined Topological Properties"

Paul Bankston

Marquette University, paul.bankston@marquette.edu

Published version. *The Journal of Symbolic Logic*, Vol. 56, No. 2 (June 1991): 425-426. [DOI](#). © 1991 Association for Symbolic Logic. Used with permission.

CORRIGENDUM TO "TAXONOMIES OF MODEL-THEORETICALLY DEFINED TOPOLOGICAL PROPERTIES"

PAUL BANKSTON

Abstract. An error has been found in the cited paper; namely, Theorem 3.1 is false.

1. I would like to correct a simple, but serious, error in [1]; namely Theorem 3.1 therein is quite false: It *can* happen that there are compact Hausdorff spaces X and Y with $X \equiv Y$ (indeed $X \cong Y$) but $X \not\equiv_{\mathbf{T}} Y$. I am most grateful to Lutz Heindorf for communicating [3] the following straightforward example: Let X and Y be any two Boolean spaces with infinite dense sets of isolated points. Then $B(X)$ and $B(Y)$ are Wallman bases for X and Y respectively, are infinite atomic Boolean algebras, and hence, by the Tarski invariants theorem, are elementarily equivalent. Thus $X \equiv Y$. However, one can easily pick X and Y as above so that $X \not\equiv_{\mathbf{T}} Y$; e.g., let X and Y be the ordinal spaces $\omega + 1$ and $\omega^2 + 1$ respectively. Then Y has a point of Cantor-Bendixson derivative 2, while X does not. This fact can be expressed in a sentence of Φ_1 .

The faulty inference in the proof of Theorem 3.1 of [1] occurs in the penultimate sentence: If W and Z are two Tichonov spaces with Wallman bases that are lattice-isomorphic, it does *not* generally follow that W and Z are homeomorphic. (We could make the inference if either W and Z were both compact or the Wallman bases contained all the singletons, but in our case W and Z are topological ultrapowers and neither condition holds.)

2. In Professor Heindorf's communication [3], there were some further interesting facts that enrich the content of [1].

2.1. The 3-cell \mathcal{S}^3 is characterized by \mathbf{T}_F in {metrizable} [2]. (This augments Theorem 1.2 in [1].)

2.2. There is a complete description of the spaces that are (finitely) characterized by certain taxonomies in {metrizable Boolean}. Let \mathcal{R} be the class of R. S. Pierce's "compact 0-dimensional metric spaces of finite type" [7].

THEOREM [5]. *For any metrizable Boolean space X , the following are equivalent:*

(i) $X \in \mathcal{R}$.

Received September 17, 1990.

1980 *Mathematics Subject Classifications*. (1985 Revision) Primary: 03C15, 03C20, 06D99, 54D30, 54F15, 54F25

(ii) X is finitely characterized by T_F in $\{\text{metrizable Boolean}\}$.

(iii) X is finitely characterized by T_i in $\{\text{metrizable Boolean}\}$. ■

(This result addresses issue (I2) in [1].)

2.3. THEOREM [6]. *There are $c T_i$ -taxa (hence $c T_F$ -taxa) in $\{\text{metrizable Boolean}\}$.* ■

(This result addresses issue (I3) in [1], and answers a question raised in the penultimate paragraph on p. 592 therein. See also the paragraphs following the proof of Theorem 2.10.)

2.4. THEOREM [4]. $\{\text{metrizable Boolean}\}$ is dense in $\{\text{Boolean}\}$, relative to T_i . ■

(This result addresses issue (I6) in [1].)

REFERENCES

- [1] P. BANKSTON, *Taxonomies of model-theoretically defined topological properties*, this JOURNAL, vol. 55 (1990), pp. 589–603.
- [2] H. G. BOTHE, *A first order characterization of 3-dimensional manifolds*, *Workshop on extended model theory* (H. Herre, editor), Report R-Math. 03/81, Akademie der Wissenschaften der DDR, Berlin, 1981, pp. 1–19.
- [3] L. HEINDORF (private communication).
- [4] ———, *Comparing the expressive power of various languages for Boolean algebras*, *Zeitschrift für Mathematische Logik und Grundlagen der Mathematik*, vol. 27 (1981), pp. 419–434.
- [5] ———, *Beiträge zur Modelltheorie der Booleschen Algebren*, Seminarbericht No. 53, Humboldt-Universität, Berlin, 1984.
- [6] B. MOLZAN, *On the number of different theories of Boolean algebras in several logics*, *Workshop on extended model theory* (H. Herre, editor), Report R-Math. 03/81, Akademie der Wissenschaften der DDR, Berlin, 1981, pp. 102–113.
- [7] R. S. PIERCE, *Compact zero-dimensional metric spaces of finite type*, *Memoirs of the American Mathematical Society*, no. 130 (1972).

DEPARTMENT OF MATHEMATICS, STATISTICS AND COMPUTER SCIENCE
MARQUETTE UNIVERSITY
MILWAUKEE, WISCONSIN 53233